



# THE CORRELATION BETWEEN THE NUMBER OF VASCULAR RISK FACTORS AND THE ONSET OF HEMODIALYSIS WITH COGNITIVE IMPAIRMENT IN CHRONIC KIDNEY DISEASE (CKD) PATIENTS

Priyanka Ganesa Utami\*, Wiwin Sundawiyani

\*Correspondence: [priyanka.utami@gmail.com](mailto:priyanka.utami@gmail.com)

Department of Neurology, Islamic Jakarta Cempaka Putih General Hospital, Jakarta, Indonesia

## Article History:

Received: October 2, 2023  
Accepted: November 15, 2023  
Published: January 1, 2024

## Cite this as:

Utami PG, Sundawiyani W. The Correlation Between the Number of Vascular Risk Factors and The Onset of Hemodialysis with Cognitive Impairment in Chronic Kidney Disease (CKD) Patients. *Magna Neurologica*. 2(1) January 2024: 11-15. 10.20961/magnaneurologica.v2i1.904

## ABSTRACT

**Background:** Cognitive impairment is a common disorder in patients with chronic kidney disease (CKD). At least, 50% of CKD patients experience impaired cognitive function caused by ischemia associated with impaired vascular function. In addition, significant changes during the dialysis process such as loss of water and ultrafiltration will disrupt the blood supply to the brain and exacerbate ischemia that underlies the incidence of impaired cognitive function.

**Objective:** This study aims to analyze the correlation between number of vascular risk factors and onset of dialysis with cognitive impairment in CKD patients

**Methods:** This research is an analytic observational study. The sample in this study was taken from chronic kidney disease patients undergoing hemodialysis treatment during February-July 2023. The sampling technique used purposive sampling. Patients who met the inclusion and exclusion criteria were interviewed and examined for cognitive function using the MOCA-INA. The statistical analysis used was Spearman's rho (non-parametric) with a 95% confidence level and  $p < 0.05$ .

**Results:** The results of correlation test using Spearman Rho showed patients undergoing hemodialysis for more than 5 years had a correlation with impaired cognitive function,  $P=0.02$ . Cognitive domains that are most affected sequentially are memory/delayed recall, visuospatial, abstraction, language, attention and naming. While the number of vascular risk factors did not have a correlation with cognitive impairment with  $P=1.38$ .

**Conclusion:** CKD patients undergoing hemodialysis for more than 5 years have a correlation with impaired cognitive function, with the most affected cognitive domains sequentially are memory or delayed recall, visuospatial, abstraction, language, attention and naming. However, the number of vascular risk factors has no correlation with cognitive impairment.

**Keywords:** chronic kidney disease, cognitive impairment, onset of hemodialysis



This is an open access article distributed under the terms of the Creative Commons Attribution- 4.0 International License

## Introduction

Chronic renal failure (CKD) is defined as a glomerular filtration rate (GFR) less than 60 mL/min/1.73 m<sup>2</sup> for more than 3 months or kidney damage resulting from structural or functional

abnormalities other than decreased GFR, which is characterized by albuminuria; accumulation of urinary sediment, electrolyte disturbances, and other abnormalities or impairment detected by histology or

imaging. End Stage Renal Disease (ESRD) occurs due to loss of kidney function with a GFR of less than 15 mL/minute requiring hemodialysis or kidney transplantation.<sup>1-3</sup> Cognitive impairment and deficits in one or more brain functions such as memory, executive function, and sensory processing are common in patients with CKD, with a prevalence that depends on the stage of CKD<sup>4</sup>. Murray et al studied 374 patients with chronic renal failure, found that 50% of them had moderate cognitive impairment and 37% had severe cognitive impairment.<sup>5-7</sup> Cognitive impairment begins early in the course of CKD and worsens as kidney function worsens. In the final stage of CKD, which requires hemodialysis, 85% of patients experience memory impairment, impaired executive function, or language deficits. Because many CKD patients suffer from impaired cognitive function, CKD is considered as a risk factor for mild cognitive impairment and dementia.<sup>4</sup>

Several risk factors that have been studied recently show that age, female gender, educational status, diabetes mellitus, hypertension, and cardiovascular disease are risk factors for impaired cognitive function in CKD patients.<sup>4,8-10</sup> Various mechanisms play a role in the process of impaired cognitive function in patients with CKD, including vascular lesions and impaired cerebral blood flow autoregulation, neuroinflammation, urea neurotoxins and renal neurotrophins. Low GFR is believed to be an independent risk factor for cognitive impairment.<sup>1,5</sup> Impaired cognitive function increases by 15-25% for every 10 ml/min decrease in GFR per 1.73 m<sup>2</sup>. This shows that patients with CKD have at least two times the risk of impaired cognitive function compared to those who do not suffer from CKD. If the GFR decreases to <30 ml/minute per 1.73 m<sup>2</sup>, this risk will increase up to four times. This is equivalent to 6-7 years of aging in cognitive function compared to the general population.<sup>3,5,6</sup> In addition, ischemic condition in the subcortical microvascular also underlie the process of cognitive function disorders in CKD patients.<sup>7-9</sup>

Cognitive impairment is also common in chronic kidney disease and in patients on dialysis. Patients undergoing hemodialysis have worse cognitive function than the general population on tests of global cognition, attention and orientation, concepts and reasoning, motor construction and performance, executive function, language, and memory.<sup>24</sup> This phenomenon is due to significant changes in the circulatory system during the dialysis process. One of them is the loss of water, both in the process of ultrafiltration and from migration to the tissues from the blood vessels which causes a decrease in blood volume and an increase in its density and viscosity, as well as an increase in peripheral resistance which causes a disruption in the blood supply of tissues, including the central nervous system.<sup>10,11</sup> Radiological

study by Mizumasa et al. also showed that the hypotension experienced during hemodialysis affected the extent of the lesion in the frontal lobe white matter. The extent of these lesions will then increase the risk of frontal lobe atrophy.<sup>12</sup>

## Methods

This research has passed the ethical review of the Health Research Ethics Committee at Islamic Jakarta Cempaka Putih General Hospital through Letter Number of 009/EC/KEP/05/2023. This research was conducted with a quantitative observational analytic approach. This research was conducted in the hemodialysis unit of the Jakarta Islamic Hospital Cempaka Putih. The sample in this study was taken from chronic kidney disease patients undergoing hemodialysis treatment during February-July 2023. The sampling technique used was purposive sampling, by selecting subjects who met the inclusion and exclusion criteria as follows:

### Inclusion Criteria

1. Patients with chronic kidney disease undergoing hemodialysis at Islamic Jakarta Cempaka Putih General Hospital
2. 30-60 years old
3. The patient is fully awake and cooperative
4. Hgb > 8 g/dL
5. Sodium > 125 mEq/L
6. Blood sugar level >60 and <400 mg/dL

### Exclusion Criteria

1. Have a history of stroke as evidenced by the results of a head CT scan
2. Have an intracranial lesion as evidenced by imaging studies
3. Having a neurological deficit based on neurological examination

The minimum sample size is calculated using the open-epi application with a design effect for cluster survey (DEEF) power of 0.75. Based on the calculation of the number of samples, with the prevalence of CKD in Indonesia based on the 2018 Riskesdas of 3.8%<sup>17</sup>, a minimum sample of 43 is obtained. Patients who met the inclusion and exclusion criteria were interviewed to meet the characteristics of the research data, number of vascular risk factors, and the onset since the first time they underwent hemodialysis. The onset of hemodialysis was divided into two groups with a cut off of 5 years: less than 5 years and more than 5 years. The patient then underwent a MOCA-INA examination with a normal value of  $\geq 26$  and an abnormal value (impaired cognitive function) of <26. Statistical analysis using Spearman Rho (95% confidence level and  $p < 0.05$ ) was then performed to see the relationship between the onset of hemodialysis with impaired cognitive function.

## Results

During April 2023, there were 168 patients diagnosed with chronic kidney failure and undergoing hemodialysis. Of the 168 patients, we sampled 45 patients who met the inclusion and exclusion criteria with the following basic characteristics (Table 1).

**Table 1.** Characteristics of Sample Data

Variables	Frequency	%
Gender		
Male	25	57%
Female	19	43%
Age		
30-50 years old	20	45%
>50 years old	24	55%
Onset of CKD		
<1 year	10	23%
1-5 years	24	55%
6-10 years	7	16%
Level of education		
Elementary School	1	2%
Junior High School	12	30%
Senior High School	20	43%
Bachelor	12	25%
MOCAINA Examination Results		
Normal	23	50%
MCI	20	45%
Cognitive Impairment	2	5%
Number of Vascular Risk Factors		
≤ 1	25	64%
>1	18	36%

Table 1 illustrates that the majority of kidney disease patients undergoing hemodialysis were aged fifty years and above, accounting for 51% of the study participants. Additionally, a significant proportion (51%) had been living with chronic kidney failure for approximately 1-5 years. Most of the patients who underwent hemodialysis and were involved in this research had an educational background of more than nine years, encompassing high school and undergraduate levels. The length of education was recognized as an influential factor and was taken into account during the MOCA-INA (Montreal Cognitive Assessment-Indonesian Version) examination. Based on the MOCA-INA results, 51% of patients were classified as having normal cognitive function, while 44% were found to have Mild Cognitive Impairment, and a small percentage (4%) exhibited signs of Cognitive Impairment. Furthermore, nearly 44% of the patients were identified as having more than one vascular risk factor. The correlation analysis results between the duration of hemodialysis and impaired cognitive function, assessed using the Spearman Rho test, are presented in Table 2.

**Table 2.** Correlation Between the Onset Hemodialysis and Impaired Cognitive Function

Variables		Cognitive Impairment
Onset of hemodialysis	Correlation Coefficient	0.367
	P value	0.07
Number of vascular risk factors	Correlation Coefficient	-0.161
	P value	0.148

Table 2 shows a significant relationship ( $P=0.02$ ;  $P<0.05$ ) between the duration of hemodialysis and impaired cognitive function. The correlation between a number of vascular risk factors and cognitive function is not significant, with P value of 0.138.

**Table 3.** Most Influenced Cognitive Domain

Domain	Mean	Normal Value	Percontation
Memory/Delayed Recall	3,4	5	68%
Visuospatial/ Executive Function	3,6	5	72%
Abstract Thinking	1,5	2	77%
Language	2,4	3	78%
Attention	4,9	6	81%
Naming	2,7	3	90%
Orientation	5,4	6	90%

## Discussion

Chronic kidney disease is a risk factor for cognitive impairment and has been proven by many studies.<sup>23</sup> Many factors cause impaired cognitive function that occurs in CKD patients. One of them is hemodialysis. Another study by Herman et al<sup>13</sup> using MMSE shows a correlation between the onset of dialysis and cognitive impairment after 12 months, while another study by Rustanti using MMSE shows no correlation between the onset of dialysis even after 24 months. Our study uses MOCAINA to assess cognitive impairment due to its sensitivity to assess mild cognitive impairment. We uses a 5-year cut-off to assess the relationship between the onset since the first time undergoing hemodialysis and impaired cognitive function, which was showed that there might be a positive correlation between the onset of hemodialysis over 5 years and cognitive impairment. This means that the longer the onset of the first time undergoing dialysis, the higher the probability of becoming cognitively impaired (Table 3).<sup>15</sup>

Cognitive impairment in dialysis patient may be related to the statement of Xie et al<sup>4</sup> which explains that cognitive impairment will worsen as kidney function worsens, since the patient who undergone dialysis has

reached end stage renal failure. Stated that the hemodialysis process and the intense catabolic characteristics of CKD may contribute to and even accelerate brain aging so that patients undergoing hemodialysis have poorer cognitive test performance.<sup>17</sup> In their study they showed that the most influential domains of impaired cognitive function were attention and calculation. Meanwhile, our research shows that the most affected cognitive domains, respectively, are memory/delayed recall, visuospatial, abstraction, language, attention, and naming. This process was described by Olczyk et al<sup>14</sup> who stated that significant changes in the circulatory system during the dialysis process would cause water loss and disruption of the brain's blood supply resulting in a decrease in white matter volume. As a result, patients undergoing hemodialysis have worse cognitive function than the general population on tests of global cognition, attention and orientation, concepts and reasoning, motor construction and performance, executive function, language, and memory.<sup>19-21</sup>

Although vascular risk factors are closely related to impaired cognitive function in CKD patients.<sup>18,22</sup> Our results show that more vascular risk factors are not correlated with impaired cognitive function, since almost all of our patients have a similar number of vascular risk factors, and most of these vascular risk factors have been controlled during their hospitalization. Even so, a lot of literature state that vascular risk factors have an important role in the occurrence of impaired cognitive function in CKD patients.<sup>4,14</sup> We realize that our study has its own limitations. Therefore, we suggest other study to assess any risk factors that might cause cognitive dysfunction in patients with chronic kidney disease. We also suggest other study to compare cognitive function before and after undergoing hemodialysis.

We realize that our study has some limitations: (1) A lot of confounding factors cannot be excluded (example: B12 and folic acid deficiency) because our facility cannot provide the laboratory test for those parameters, (2) Further studies are needed to really conclude the correlation between dialysis and cognitive impairment.<sup>24,25</sup> For example: comparison of cognitive impairment between dialysis and dialysis patient or comparison between two different type dialysis duration (fast/slow).

## Conclusion

Our study concludes that CKD patients undergoing hemodialysis for more than 5 years have a positive correlation with impaired cognitive function, means longer the patient had undergone dialysis will probably have the higher chance to had cognitive impairment. The most affected cognitive domains sequentially are

memory or delayed recall, visuospatial, abstraction, language, attention and naming. However, the number of vascular risk factors has no correlation with cognitive impairment. Further study to compare groups with and without dialysis or imaging study should be reconsidered.

## Declaration of Interests

We have no relevant financial interest to disclose.

## References

1. Li X, Yuan F, Zhou L. Organ Crosstalk in Acute Kidney Injury: Evidence and Mechanisms. *J Clin Med*; 2022 Nov. 11(22):6637. DOI: 10.3390/jcm11226637
2. Grasing M, Sharma P, Lepping RJ, Honea R, Burns JM, Brooks WM, et al. Association between the Estimated Glomerular Filtration Rate and Brain Atrophy in Older Adults. *Am J Nephrol*; 2022. 53(2–3):176–81. DOI: 10.1159/000521892
3. Liu HS, Hartung EA, Jawad AF, Ware JB, Laney N, Port AM, et al. Regional cerebral blood flow in children and young adults with chronic kidney disease. *Radiology*; 2018. 288(3):849–58. DOI: 10.1148/radiol.2018171339
4. Xie Z, Tong S, Chu X, Feng T, Geng M. Chronic Kidney Disease and Cognitive Impairment: The Kidney-Brain Axis. *Kidney Dis*; 2022. 8(4):275–85. DOI: 10.1159/000524475
5. Bikbov B, Soler MJ, Pešić V, Capasso G, Unwin R, Endres M, et al. Albuminuria as a risk factor for mild cognitive impairment and dementia-what is the evidence?. *Nephrol Dial Transplant*; 2021. 37:ii55–62. DOI: 10.1093/ndt/gfab261
6. Bronas UG, Puzantian H, Hannan M. Cognitive impairment in chronic kidney disease: Vascular milieu and the potential therapeutic role of exercise. *Biomed Res Int*; 2017. 2017: 1273150. DOI: 10.1155/2017/1273150
7. Lau WL, Huisa BN, Fisher M. The Cerebrovascular-Chronic Kidney Disease Connection: Perspectives and Mechanisms. *Transl Stroke Res*; 2017. 8(1): 67–76. DOI: 10.1007/s12975-016-0499-x
8. Drew DA, Weiner DE, Sarnak MJ. Cognitive Impairment in CKD: Pathophysiology, Management, and Prevention. *Am J Kidney Dis*; 2019. 74(6):782–90. Retrieved on Mar 14, 2023. Available from: <https://doi.org/10.1053/j.ajkd.2019.05.017>
9. Long Y, Nie J. Homocysteine in Renal Injury. *Kidney Dis*; 2016. 2(2):80–7. DOI: 10.1159/000444900
10. Olczyk P, Kusztal M, Gołębiowski T, Letachowicz K, Krajewska M. Cognitive Impairment in End Stage Renal Disease Patients Undergoing Hemodialysis:

- Markers and Risk Factors. *Int J Environ Res Public Health*; 2022. 19:4. DOI: 10.3390/ijerph19042389
11. Gäckler A, Rohn H, Lisman T, Benkö T, Witzke O, et al. Evaluation of hemostasis in patients with end-stage renal disease. *PLoS One*; 2019. 14 (2):1–13. DOI: 10.1371/journal.pone.0212237
  12. Tsuruya K, Yoshida H. Cognitive Impairment and Brain Atrophy in Patients with Chronic Kidney Disease. *J Clin Med*; 2024 Feb. 28;13(5):1401. DOI: 10.3390/jcm13051401
  13. Herman I, Yonata A, Tjiptaningrum A, Berawi KN, Kedokteran F, Lampung U, et al. Hubungan Lama Hemodialisis dengan Fungsi Kognitif Pasien Penyakit Ginjal Kronik yang Menjalani Hemodialisis di RSUD Abdul Moeloek Bandar Lampung The Relationship between Duration of Hemodialysis and Cognitive Function Chronic Kidney Disease Patient in Hem. *Medula*; 2017. 7(5) :47–53. Retrieved on Mar 12, 2023. Available from: <https://academicjournal.yarsi.ac.id/index.php/sainsteke/s/article/view/1606>
  14. Miglinas M, Cesniene U, Janusaite MM, Vinikovas A. Cerebrovascular Disease and Cognition in Chronic Kidney Disease Patients. *Front Cardiovasc Med*; 2020. 7 (June):1–13. DOI: 10.3389/fcvm.2020.00096
  15. Freitas ME, Ruiz-Lopez M, Dalmau J, Erro R, Privitera M, Andrade D, et al. Seizure and Movement Disorders: Phenomenology, Diagnostic Challenges, and Therapeutic Approaches. *J Neurol Neurosurg Psychiatry*; 2019. 90: 920-928. DOI: 10.1136/jnnp-2018-320039
  16. Hidayat R, Widjaya N, Djuliannisaa Z, et al. Ischemic stroke in COVID-19 patients: a cross-sectional study from an Indonesian COVID-19 referral hospital. *Egypt J Neurol Psychiatry Neurosurg*; 2022. 58(1). DOI: 10.1186/s41983-022-00528-z
  17. Weitz JI, Fredenburgh JC, Eikelboom JW. A Test in Context: D-Dimer. *J Am Coll Cardiol*; 2017. 70(19):2411-2420. DOI: 10.1016/j.jacc.2017.09.024
  18. Yao Y, Cao J, Wang Q, et al. D-dimer as a biomarker for disease severity and mortality in COVID-19 patients: A case control study. *J Intensive Care*; 2020. 8(1):1-11. DOI: 10.1186/s40560-020-00466-z
  19. Widjaja AC. Uji diagnostik pemeriksaan kadar d-dimer plasma pada diagnosis stroke iskemik. [Tesis]. Universitas Diponegoro; 2014. p. 29-30. Retrieved on Mar 14, 2023. Available from: [https://eprints.undip.ac.id/24038/?utm\\_source](https://eprints.undip.ac.id/24038/?utm_source)
  20. Kuriakose D, Xiao Z. Pathophysiology and treatment of stroke: Present status and future perspectives. *Int J Mol Sci*; 2020. 21(20):1-24. DOI: 10.3390/ijms21207609
  21. Atmaja KS, Wicaksana AAGOS, Putra IWAS, Putra WWS. Hubungan konsentrasi serum C-Reactive Protein dan D-dimer dengan derajat keparahan dan mortalitas pasien COVID-19. *Intisari Sains Medis*; 2021. 12(2):680-685. DOI: 10.15562/ism.v12i2.971
  22. Mus R, Thaslifa T, Abbas M, Sunaidi Y. Studi Literatur: Tinjauan Pemeriksaan Laboratorium pada Pasien COVID-19. *J Kesehat Vokasional*; 2021. 5(4):242. DOI: 10.22146/jkesvo.58741
  23. Sabatudung C, Bia MB, Kambuno NT. Gambaran Hematologi Rutin Dan D-Dimer Pasien COVID-19 Kota Kupang. *J Indones Med Lab Sci*; 2023. 4(1):14-24. DOI: 10.53699/joimedlabs.v4i1.125
  24. Zhang L, Sun W, Wang Y, et al. Clinical Course and Mortality of Stroke Patients With Coronavirus Disease 2019 in Wuhan, China. *Stroke*; 2020. 51(9):2674-2682. DOI: 10.1161/STROKEAHA.120.030642
  25. Akbar NA, Rahardjo AM, Parti DD, Sakinah EN. Analysis of the Relationship between Laboratory Biomarkers and the Severity of COVID-19 at Kaliwates General Hospital Jember. *J Agromedicine Med Sci*; 2022. 8(1):51. DOI: 10.19184/ams.v8i1.29062